

On rational points of orthogonal group

Masanori Kobayashi, Chikara Nakayama

Abstract

Let n be a positive integer. We show that a unit rational space vector whose multiple by n is an integer vector can be extended to a rational orthonormal basis whose all members have the same property.

1 Main results

Definition 1.1 A system $S : v_1, \dots, v_k$ of vectors is called **orthoregular** if (i) v_i and v_j are perpendicular to each other if $i \neq j$, and (ii) $|v_i| = |v_j| \neq 0$. We call $|v_i|$ the **length** of the system.

Proposition 1.2 If n is odd, the length of orthoregular n vectors in $\mathbf{Z}^n$ is an integer.

Proof. Let $v_1, \dots, v_n$ be orthoregular integer vectors and l be its length. Then the determinant $|v_1, \dots, v_n| = |v_1| \cdots |v_n| = l^n$ is an integer since all the entries are integers. Since l^2 is an integer, we have $l \in \mathbf{Q}$, and $l \in \mathbf{Z}$ by the normality of $\mathbf{Z}$.

Definition 1.3 Let k be a positive integer less than n . We denote the maximum number of orthoregular integer vectors extending an orthoregular system $S \subset \mathbf{Z}^n$ by $E(S)$, and the minimum of those $E(S)$ for all orthoregular S with $\sharp S = k$ by $E(n, k)$. For a subset K of $\mathbf{R}$, we also use $E_K(n, k)$ for all orthoregular sets of k integer vectors of length in K . We mainly use the cases $K = \mathbf{R}$ or $\mathbf{Z}$.

Proposition 1.4 If n is even or $K = \mathbf{Z}$, $E_K(n, n-1) = n$.

Proof. Let $v_1, \dots, v_{n-1}$ be an orthoregular system in $\mathbf{Z}^n$ of length l . Take a square matrix $(v_1, \dots, v_{n-1}, v)$, where v is an arbitrary vector. Let v_n be the vector consists of the cofactors of the last column. Then v_n is orthogonal to $v_1, \dots, v_{n-1}$, thus $l^{n-1}|v_n| = |v_n|^2$ by cofactor expansion. Hence $|v_n| = l^{n-1}$.

We will now show that $(1/l^{n-2})v_n \in \mathbf{Z}^n$, which will complete the proof. It is enough to show that the last component of v_n is in $l^{n-2}\mathbf{Z}$ since the others are shown similarly. We put $v_j = \begin{pmatrix} v'_j \\ x_j \end{pmatrix}$ for $1 \leq j \leq n-1$, where v'_j has $(n-1)$ -rows and x_j is a scalar. We show that $D := |v'_1, \dots, v'_{n-1}|$ belongs to $l^{n-2}\mathbf{Z}$. Since $l^{n-2} \in \mathbf{Z}$ by the assumption that n is even or $l \in \mathbf{Z}$, it is enough to prove that $D^2 \in l^{2(n-2)}\mathbf{Z}$.

$$D^2 = |(v'_i, v'_j)_{1 \leq i, j \leq n-1}| = |((v_i, v_j) - x_i x_j)_{i, j}|,$$

which is nothing but $\varphi(l^2)$, where $\varphi(t)$ is the characteristic polynomial for the matrix $(x_i x_j)_{i, j}$. Since the matrix has rank ≤ 1 , $\varphi(t)$ has a form $t^{n-2}(t - a)$ ($a \in \mathbf{Z}$). Thus $D^2 \in l^{2(n-2)}\mathbf{Z}$.

The next is the main result in this paper.

Theorem 1.5 *A nonzero integer vector v can be extended to an integral orthoregular basis if and only if $|v|$ is an integer. In particular, $E_{\mathbf{Z}}(3, 1) = 3$.*

Examples. Here is the list of smallest primitive nonnegative integral space vectors whose norms are integers.

${}^t(0, 0, 1)$, ${}^t(1, 2, 2)$, ${}^t(0, 3, 4)$, ${}^t(2, 3, 6)$, ${}^t(1, 4, 8)$, ${}^t(4, 4, 7)$, ${}^t(2, 6, 9)$, ${}^t(6, 6, 7)$, ...

By starting from each vector in the list, you can always complete it to an orthoregular basis consisting of integral vectors. Dividing by its norm, you get a rational orthonormal basis, thus a rational point of $O(3)$.

2 Proof of main theorem

In this section, we prove 1.5. Our proof is constructive and gives an algorithm for all solutions.

First we prove two lemmas.

Lemma 2.1 *Let $v, w \in \mathbf{Z}^3 \setminus \{\mathbf{o}\}$. Assume $|v| \in \mathbf{Z}$ and $(v, w) = 0$. Then $v_p(|w|^2)$ is even for any prime p with $p \equiv 3 \pmod{4}$. Here $(-, -)$ is the standard inner product and v_p is the normalized valuation with respect to p .*

Proof. Let $v = \begin{pmatrix} a \\ b \\ c \end{pmatrix}$, $w = \begin{pmatrix} x \\ y \\ z \end{pmatrix}$ and let $l = |v|$. We may assume $a \neq 0$. Since $ax + by + cz = 0$, we have $x = \frac{-by - cz}{a}$. Hence

$$|w|^2 = \left(\frac{by + cz}{a} \right)^2 + y^2 + z^2,$$

$$a^2|w|^2 = (a^2 + b^2)y^2 + 2bcyz + (a^2 + c^2)z^2.$$

Multiplying $a^2 + b^2$, we deduce

$$a^2(a^2 + b^2)|w|^2 = ((a^2 + b^2)y + bcz)^2 + ((a^2 + c^2)(a^2 + b^2) - b^2c^2)z^2.$$

But the coefficient of z^2 in the right-hand-side coincides with a^2l^2 , so that the both terms of the right-hand-side are perfect squares. Hence v_p of the right-hand-side is even for any p with $p \equiv 3 \pmod{4}$. Since $v_p(a^2(a^2 + b^2))$ is also even, $v_p(|w|^2)$ is even, too.

Lemma 2.2 *Let $a, b, c, l \in \mathbf{Z}$ satisfying $ac - b^2 = l^2$. Then there exist $x, y \in \mathbf{Z}$ such that $ax^2 + 2bxy + cy^2 = l^2$ if and only if $a \geq 0$ and, when $a > 0$, $v_p(a)$ is even for any prime p with $p \equiv 3 \pmod{4}$.*

Proof. It is probably well-known. Our proof here uses the prime factorization in the integer ring $\mathbf{Z}[i]$, where $i = \sqrt{-1}$.

We prove the only if part. By multiplying a with the given equation, we have $(ax + by)^2 + (ly)^2 = al^2$. Hence l divides $ax + by$. Letting $u = \frac{ax + by}{l}$, we have $u^2 + y^2 = a$, which implies that the desired condition holds.

We prove the if part. The case $a = 0$ is trivial. Assume $a > 0$. If there are u and y with $u^2 + y^2 = a$ such that a divides $ul - by$, then $x = \frac{ul - by}{a}$ and y give an integer solution for

$ax^2 + 2bxy + cy^2 = l^2$. We prove the existence of u and y as above under the assumption that $v_p(a)$ is even for any prime p with $p \equiv 3 \pmod{4}$. First factorize a into

$$a = 2^m \prod_{p_j \equiv 1 \pmod{4}} p_j^{m_j} \prod_{q_k \equiv 3 \pmod{4}} q_k^{n_k},$$

where p_j 's are primes with $p_j \equiv 1 \pmod{4}$ and q_k 's are primes with $q_k \equiv 3 \pmod{4}$. Here, each n_k is even by the assumption. Since $l^2 + b^2 = ac$, we can write $l + ib$ as a product

$$l + ib = u_0(1 + i)^{m'} \prod p_j^{s_j} \pi_j^{t_j} \prod q_k^{n'_k/2}$$

in $\mathbf{Z}[i]$, where u_0 is a unit of $\mathbf{Z}[i]$, $m' \geq m$, $p_j = \pi_j \bar{\pi}_j$ in $\mathbf{Z}[i]$, where π_j and $\bar{\pi}_j$ are conjugate prime elements, $2s_j + t_j \geq m_j$, and $n'_k \geq n_k$ is an even.

Claim. There exist $a_j \geq 0, b_j \geq 0$ such that $s_j + \min(t_j, b_j) \geq a_j + b_j$ and $2a_j + b_j = m_j$.

Proof of Claim. If $2s_j \leq m_j$, let $a_j = s_j$ and $b_j = m_j - 2s_j$. Then, we have $t_j \geq m_j - 2s_j = b_j$ and a_j, b_j satisfy the condition.

If $2s_j > m_j$, let $a_j := \lfloor \frac{m_j}{2} \rfloor$ and $b_j = m_j - 2\lfloor \frac{m_j}{2} \rfloor$. Then, we have $b_j \leq 1$ and $s_j \geq a_j + 1 \geq a_j + b_j$ and a_j, b_j satisfy the condition. This completes the proof of Claim.

Using these a_j and b_j , define

$$u + iy := (1 + i)^m \prod p_j^{a_j} \bar{\pi}_j^{b_j} \prod q_k^{n_k/2}.$$

Then, since $2a_j + b_j = m_j$, we have $u^2 + y^2 = a$. Further, $ul - by$, the real part of $(u + iy)(l + ib)$, is divided by $2^m \prod p_j^{a_j + s_j + \min(b_j, t_j)} \prod q_k^{n_k}$. Since $a_j + s_j + \min(b_j, t_j) \geq 2a_j + b_j = m_j$, this is divided by a .

Now we prove 1.5. The only if part is by 1.2. We prove the if part. By 1.4, it is enough to show $E_{\mathbf{Z}}(3, 1) \geq 2$. Let $v_1 = \begin{pmatrix} a_1 \\ b_1 \\ c_1 \end{pmatrix} \in \mathbf{Z}^3 \setminus \{\mathbf{o}\}$ such that $l := |v_1|$ is an integer. It suffices to show $E_{\mathbf{Z}}(\{v_1\}) \geq 2$. Since $m := (a_1, b_1, c_1)$ divides l , replacing v_1 with $\frac{v_1}{m}$, we may and will assume that $m = 1$. Then, the homomorphism $(v_1, -): \mathbf{Z}^3 \rightarrow \mathbf{Z}$ is surjective. The kernel K of this surjection is free of rank two. Let Q be the positive definite bilinear form on K obtained by restricting the standard metric on $\mathbf{Z}^3$. It is enough to show that Q represents l^2 . Take a base w_1, w_2 of K . Let $A := \text{Gram}(w_1, w_2)$, the symmetric matrix which represents Q with respect to w_1, w_2 .

Claim. $\det(A) = l^2$.

Proof of Claim. Let S be the area of the parallelogram spanned by w_1, w_2 . What we have to see is $S = l$. Since there are isomorphisms $\mathbf{Z}^3 / \langle v_1, w_1, w_2 \rangle \cong \mathbf{Z} / (v_1, v_1)\mathbf{Z} \cong \mathbf{Z} / l^2 \mathbf{Z}$, the volume of the parallelepiped spanned by v_1, w_1, w_2 is l^2 . On the other hand, since v_1 is perpendicular to K , this volume is $|v_1|S$, too. Hence $S = l$. This completes the proof of Claim.

Let $a := (w_1, w_1)$. By 2.1 with $v = v_1$ and $w = w_1$, $v_p(a)$ is even for any prime p with $p \equiv 3 \pmod{4}$. Hence by 2.2 with $a = (w_1, w_1)$, $b = (w_1, w_2)$, $c = (w_1, w_2)$, we conclude that Q represents l^2 , which completes the proof of 1.5.

3 More results

Here we gather more results on $E_K(n, k)$.

Proposition 3.1 *For $0 < k < n$ and let K be a subset of $\mathbf{R}$.*

1. $E_{\mathbf{Z}}(n+1, k+1) \leq E_{\mathbf{Z}}(n, k) + 1$.
2. If n is odd, then $E(n, k) \leq n - 1$.
3. If n is odd and each coordinate of $v \in \mathbf{Z}^n$ is odd, then $E(v) = 1$.
4. If n is odd, then $E(n, 1) = 1$.
5. If $n \equiv 1 \pmod{8}$, then $E_{\mathbf{Z}}(n, 1) = 1$.
6. If n is the sum of positive integers n_i , $E_K(n, 1) \geq \min_i E(n_i, 1)$.
7. For a positive integer m , $E_K(mn, 1) \geq E(n, 1)$, $E_K(2m, 1) \geq 2$, $E_K(4m, 1) \geq 4$ and $E_K(8m, 1) \geq 8$.

Question. Is $E_K(16m, 1) \geq 16$ valid?

Proof. (1) Take an orthoregular system $S : v_1, \dots, v_k$ of vectors which attains $E_{\mathbf{Z}}(n, k) = E(S)$. Let l be the length of S . Then $E\left(\begin{pmatrix} l \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ v_1 \end{pmatrix}, \dots, \begin{pmatrix} 0 \\ v_k \end{pmatrix}\right) = E(v_1, \dots, v_k) + 1$.

(2) It is enough to show that $E(v_1) \leq n - 1$ for $v_1 = {}^t(1 \ 1 \ 0 \ \dots \ 0) \in \mathbf{Z}^n$. This follows from 1.2. (In fact, $E(v_1) = n - 1$. This is seen by taking $v_2 = {}^t(1 \ -1 \ 0 \ \dots \ 0), \dots, v_{2k-1} = {}^t(0 \ \dots \ 0 \ 1 \ 1 \ 0 \ \dots \ 0), v_{2k} = {}^t(0 \ \dots \ 0 \ 1 \ -1 \ 0 \ \dots \ 0), \dots$ ($1 \leq k \leq (n-1)/2$). Here the i -th components of v_{2k-1} and of v_{2k} are zero unless $i = 2k - 1, 2k$.)

(3) Suppose that an integer vector $v' = {}^t(a_1 \ \dots \ a_n)$ satisfies $(v, v') = 0$ and $|v| = |v'|$. By taking modulo two, we have $|v|^2 = |v'|^2 = a_1^2 + \dots + a_n^2 \equiv a_1 + \dots + a_n$. Since the components of v are odd, the last integer is congruent to $(v, v') = 0$, that is, $|v|^2 \equiv 0 \pmod{2}$. On the other hand, we have $|v|^2 \equiv n \equiv 1 \pmod{2}$. A contradiction.

(4) Take $v = {}^t(1 \ \dots \ 1)$ and apply (3).

(5) Use (3) and the following Claim. (The part (b) $\Rightarrow$ (a) suffices.)

Claim. For an odd integer $n > 0$, the followings are equivalent.

- (a) There exist n odd integers whose square sum is again square.
- (b) $n \equiv 1 \pmod{8}$.

Proof of Claim. (a) $\Rightarrow$ (b). Assume that there are $x_1, \dots, x_n, y \in \mathbf{Z}$ such that $x_1^2 + \dots + x_n^2 = y^2$. Since the square of any odd number is congruent to 1 modulo 8, we have $n \equiv x_1^2 + \dots + x_n^2 = y^2 \pmod{8}$. Since y is also odd, this implies $n \equiv 1 \pmod{8}$.

(b) $\Rightarrow$ (a). Take $m \geq 1$ such that $(2m-1)^2 \leq n \leq (2m+1)^2$. If we prove $n \geq \frac{(2m+1)^2 - n}{8}$, then, $(x_1, \dots, x_n) = (3, \dots, 3, 1, \dots, 1)$ will do, where the number of 3 is $\frac{(2m+1)^2 - n}{8}$, for $x_1^2 + \dots + x_n^2 = n + (9-1)\frac{(2m+1)^2 - n}{8} = (2m+1)^2$.

We prove the above inequality. This is equivalent to $9n \geq (2m+1)^2$, which reduces to $9(2m-1)^2 \geq (2m+1)^2$, $3(2m-1) \geq 2m+1$, $m \geq 1$, which completes the proof of Claim.

(6) Let $v \in \mathbf{Z}^n \setminus \{\mathbf{o}\}$. By the isomorphism $\mathbf{Z}^n \cong \mathbf{Z}^{n_1} \oplus \mathbf{Z}^{n_2} \oplus \dots$, we regard v as a collection $v_1, v_2, \dots$ with $v_i \in \mathbf{Z}^{n_i}$. Let $m := \min_i E(n_i, 1)$. For each i , if v_i is not zero, apply $E(n_i, 1)$ to v_i , and we get an orthoregular set $v_{i1} = v_i, v_{i2}, \dots, v_{im}$. If v_i is zero, let $v_{i1} = v_{i2} = \dots = v_{im}$

be zero. For each $j = 2, \dots, m$, let $v_j \in \mathbf{Z}^n$ be the vector corresponding to $(v_{ij})_i$. Then, $v_1, v_2, \dots, v_m$ is the desired system.

(7) By (6), we may assume that $m = 1$. Then this is by the next proposition with $K = \mathbf{R}$.

Proposition 3.2 *Let K be an arbitrary subset of $\mathbf{R}$ satisfying $-K = K$. If $n = 2, 4$ or 8 , then $E_K(n, 1) = n$ for any K .*

Proof. Let $v_1 \in K^n$. Via a standard matrix representation of complex number field, quaternion algebra or Cayley algebra as an $\mathbf{R}$ -algebra, the rotations multiplied by the unit imaginary numbers gives the desired $v_2, \dots, v_n$. Explicitly, for $v_1 = \begin{pmatrix} a_0 \\ a_1 \end{pmatrix}$, we give $v_2 = iv_1 = \begin{pmatrix} -a_1 \\ a_0 \end{pmatrix}$;

for $v_1 = \begin{pmatrix} a_0 \\ a_1 \\ a_2 \\ a_3 \end{pmatrix}$, we give $(v_2, v_3, v_4) = (iv_1, jv_1, kv_1) = \left(\begin{pmatrix} -a_1 \\ a_0 \\ -a_3 \\ a_2 \end{pmatrix}, \begin{pmatrix} -a_2 \\ a_3 \\ a_0 \\ -a_1 \end{pmatrix}, \begin{pmatrix} -a_3 \\ -a_2 \\ a_1 \\ a_0 \end{pmatrix} \right)$; for

$v_1 = \begin{pmatrix} a_0 \\ a_1 \\ a_2 \\ a_3 \\ a_4 \\ a_5 \\ a_6 \\ a_7 \end{pmatrix}$, we give

$$(v_2, v_3, v_4, v_5, v_6, v_7, v_8) = \left(\begin{pmatrix} -a_1 \\ a_0 \\ -a_3 \\ a_2 \\ -a_5 \\ a_4 \\ a_7 \\ -a_6 \end{pmatrix}, \begin{pmatrix} -a_2 \\ a_3 \\ a_0 \\ -a_1 \\ -a_6 \\ -a_7 \\ a_4 \\ a_5 \end{pmatrix}, \begin{pmatrix} -a_3 \\ -a_2 \\ a_1 \\ a_0 \\ -a_7 \\ a_6 \\ -a_5 \\ a_4 \end{pmatrix}, \begin{pmatrix} -a_4 \\ a_5 \\ a_6 \\ a_7 \\ a_0 \\ -a_1 \\ -a_2 \\ -a_3 \end{pmatrix}, \begin{pmatrix} -a_5 \\ -a_4 \\ -a_6 \\ a_7 \\ a_1 \\ a_0 \\ a_3 \\ -a_2 \end{pmatrix}, \begin{pmatrix} -a_6 \\ -a_7 \\ -a_4 \\ a_5 \\ a_2 \\ -a_3 \\ a_0 \\ a_1 \end{pmatrix}, \begin{pmatrix} -a_7 \\ a_6 \\ -a_5 \\ -a_4 \\ a_3 \\ a_2 \\ -a_1 \\ a_0 \end{pmatrix} \right).$$

Proposition 3.3 $E(6, k) \leq 4$ for $1 \leq k \leq 4$.

Proof. Let $v_1 = {}^t(1, 1, 1, 1, 1, 1)$. It is enough to show $E(6, v_1) \leq 4$. First observe that any integer vector who has the same length as v_1 satisfies either (a) the absolute value of any its component is 1 or (b) the absolute values of its components are 2, 1, 1, 0, 0, 0.

If an integer vector satisfying (a) is perpendicular to v_1 , then it has exactly 3 components whose value is -1 . From this, we see that there is no orthoregular set v_1, v_2, v_3 such that both v_2 and v_3 satisfy (a).

On the other hand, if an integer vector satisfying (b) is perpendicular to v_1 , then it satisfies (b') its components are 2, -1 , -1 , 0, 0, 0 or -2 , 1, 1, 0, 0, 0. Further, if both v_2 and v_3 satisfy (b') and they are perpendicular, their supports should be disjoint. From this, we see that there is no orthoregular set v_1, v_2, v_3, v_4 such that three of v_2, v_3 , and v_4 satisfy (b).

Therefore there are no orthoregular set v_1, v_2, v_3, v_4, v_5 , that is, $E(6, v_1) \leq 4$, as desired.

Remark: In fact, $E(6, v_1) = 4$ as seen by $v_2 = {}^t(1, 1, 1, -1, -1, -1)$, $v_3 = {}^t(2, -1, -1, 0, 0, 0)$, $v_4 = {}^t(0, 0, 0, 2, -1, -1)$.

Masanori Kobayashi
Department of Mathematics and Information Sciences
Tokyo Metropolitan University
1-1 Minami-Ohsawa, Hachioji, Tokyo, 192-0397
Japan
kobayashi-masanori@tmu.ac.jp

Chikara Nakayama
Department of Economics
Hitotsubashi University
2-1 Naka, Kunitachi, Tokyo 186-8601
Japan
c.nakayama@r.hit-u.ac.jp